



## **Surfactants as enhancement of In Situ Alkaline Hydrolysis (ISAH) of pesticide DNAPL**

### *Results of bench and pilot scale field tests*

Muff, Jens; Søgaard, Erik Gydesen; Bennedsen, Lars R.; Rügge, Kirsten; MacKinnon, Leah ; Durant, Neal; Pennell, Kurt ; Bondgaard, Morten

*Publication date:*  
2015

*Document Version*  
Accepted author manuscript, peer reviewed version

[Link to publication from Aalborg University](#)

#### *Citation for published version (APA):*

Muff, J., Søgaard, E. G., Bennedsen, L. R., Rügge, K., MacKinnon, L., Durant, N., Pennell, K., & Bondgaard, M. (2015). *Surfactants as enhancement of In Situ Alkaline Hydrolysis (ISAH) of pesticide DNAPL: Results of bench and pilot scale field tests*. Poster presented at AquaConSoil 2015, København, Denmark.

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

#### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



# Surfactants as enhancement of *In Situ* Alkaline Hydrolysis (ISAH) of Pesticide DNAPL: Results of bench and pilot scale field tests

Jens Muff<sup>\*1</sup> and Erik G. Søgaard<sup>1</sup>, Lars Bennedsen<sup>2</sup>, Kirsten Rügge<sup>3</sup>, Leah MacKinnon and Neal D. Durant<sup>4</sup>, Kurt Pennell<sup>5</sup>, Morten Bondgaard<sup>6</sup>

<sup>1</sup>Aalborg University, Denmark, <sup>2</sup>Rambøll, Denmark, <sup>3</sup>COWI, Denmark, <sup>4</sup>Geosyntec Consultants, CA & Maryland USA, <sup>5</sup>Tufts University, Massachusetts, USA, <sup>6</sup>Central Region, Denmark

## Background

Groyne 42 is a 20,000 m<sup>2</sup> former chemical dump site in Denmark contaminated with 100-200 tons of **organophosphorous pesticides (OPPs)**. The majority of contaminant mass is present as sorbed phase and residual **DNAPL**. The **NorthPestClean** project was established to determine the effectiveness of using **in situ alkaline hydrolysis** to treat the DNAPL (Fig. 1).




Fig. 1: The alkaline hydrolysis of OPPs.

The primary challenge was *in situ* mixing, establishing sufficient contact between hydroxide and DNAPL and surfactants were tested as 1 of 3 enhancement technologies at bench and pilot scale.

## Site and sampling



Fig. 2: The site is located directly at the waterfront. Residual OPP DNAPL is widespread in hot spots.



Fig. 3: Sampling of site water and soil with visible DNAPL. Over 2½ years, more than 2000 water samples and 1200 soil samples were analyzed in NorthPestClean.

## Initial selection

Previous work has identified **non-ionic surfactants** as the best performing type of surfactants for increasing OPP solubility. Ten non-ionic candidates were tested in bench scale, and equilibrium solubility tests showed that **alcohol ethoxylate non-ionic surfactants** was superior with respect to increasing OPP solubility compared to the alkaline tap water reference.

## Data representation

Parameter	Compounds
SUM OPPs	parathion (EP3), methyl-parathion (MP3), malathion, ethyl-sulfotep
SUM P2 acids	O,O-diethylthiophosphoric acid (EP2), O,O-dimethylthiophosphoric acid (MP2)
PNP	para-nitrophenol

## Bench scale testing




Fig. 4: Batch equilibrium tests of ethoxylate surfactants at pH 13 with a 10:1 surfactant:DNAPL volumetric ratio. Reaction time was 7 days.





Fig. 5: Batch soil tests with contaminated soil / surfactant water ratio of 4 g/mL. No extra DNAPL was added. Results presented are aqueous concentrations and soil removal of four alcohol ethoxylates after 7 days of reaction at pH 13.

Ecosurf EH-9 was chosen for pilot scale testing due to best performance in soil test (lower soil sorption compared to Rhodasurf) and cost of the chemical (3-4 € / kg).



## Pilot scale testing




Fig. 6: Pilot testing was completed in 10x10 m test cells (TCs) with TC2 used for testing of surfactants in cycle 3. 1600 kg Ecosurf EH-9 was mixed with extracted water in a 25% solution and added the 60 m<sup>3</sup> effective pore volume of the targeted treatment area (TTA) resulting in a concentration of about 2.7%.

### Baseline ISAH




Fig. 7: Results showed an increase in EP3 and a slight increase in hydrolysis products up to 1 year after surfactant addition.




Fig. 8: Surface tension analysis showed an equal distribution of surfactant in the TTA of TC2.

## Conclusion

Surfactant enhanced ISAH increased dissolved OPP concentration (x10) and total mass removal was 20-40% higher compared to baseline ISAH.